

Bursting Bubbles: from Champagne to Mudpots

Description

Interaction of gas bubbles with the free liquid-gas interface is ubiquitous. For example, transporting aromatics from champagne and pathogens from contaminated water. Furthermore, the process is also responsible for the sea spray formation as a consequence of ejecting myriads of droplets.

First, the air bubble, generated in the liquid bulk, being lighter than the surrounding medium, rises because of buoyancy and reaches the liquid-air interface. It stays there as the thin film between the bubble and the free surface drains and ruptures to form film droplets (Lhuissier and Villermaux, 2012). The rupture results in the formation of a hole in the liquid meniscus. The unstable open cavity collapses leading to the interaction of the capillary waves and forms of an upward jet. Figure 1 illustrates a typical temporal sequence of the process. The phenomenon has been widely studied by Deike et al., 2018; Duchemin et al., 2002; Gañán-Calvo, 2018. Deike et al., 2018 has provided quantitative cross-validation of the numerical and experimental studies (Figure 2). Nonetheless, the literature lacks a comprehensive study on the influence of liquid pool properties on the bubble bursting process. Notably, the influence of rheological properties on the scaling laws can provide a close-up to the understanding of the phenomenon. This latter is highly related to some geophysical phenomena such as mudpots (see: <https://www.youtube.com/watch?v=a9hUsVq9q7U>).




Figure 1: Time resolution of the process. The color shows the magnitude of non-dimensionalized vorticity, Γ (maximum $\Gamma = 150$ with red and minimum $\Gamma = -150$ with blue).




Figure 2: Collapse of the bubble cavity and interaction of capillary waves

What you will do and what you will learn?

1. You will learn about the fundamental fluid mechanics of two-phase flows.
2. You will learn the science of rheology.
3. You will learn the freeware code [Basilisk C](#) to simulate fluid dynamics problems.

If you have any questions, feel to contact [Vatsal](#) or [Mazi](#) (details below).

Supervision	E-mail	Office
Vatsal Sanjay	v.sanjay@utwente.nl	Meander 212
Dr. Maziyar (Mazi) Jalaal	m.jalaal@utwente.nl	Meander 208
Prof. D. Lohse	d.lohse@utwente.nl	Meander 261

References

- Deike, Luc, Elisabeth Ghabache, Gérard Liger-Belair, Arup K Das, Stéphane Zaleski, Stéphane Popinet, and Thomas Séon (2018). “Dynamics of jets produced by bursting bubbles”. In: *Physical Review Fluids* 3.1, p. 013603.
- Duchemin, Laurent, Stéphane Popinet, Christophe Josserand, and Stéphane Zaleski (2002). “Jet formation in bubbles bursting at a free surface”. In: *Physics of fluids* 14.9, pp. 3000–3008.
- Gañán-Calvo, Alfonso M (2018). “Scaling laws of top jet drop size and speed from bubble bursting including gravity and inviscid limit”. In: *Phys. Rev. Fluids* 3 (9), p. 091601. DOI: [10.1103/PhysRevFluids.3.091601](https://doi.org/10.1103/PhysRevFluids.3.091601).
- Lhuissier, Henri and Emmanuel Villermaux (2012). “Bursting bubble aerosols”. In: *Journal of Fluid Mechanics* 696, pp. 5–44.